MiraLAX & Gatorade Prep

Purchase the Following Products:

- 1. Two (2) Dulcolax Tablets or Liquid Tablets (bisacodyl laxative)
- 2. Two (2) MiraLAX 238-gram bottles or generic equivalent
- **3. Two (2) Gatorade 64-ounce bottles (NOTHING RED, ORANGE OR PURPLE).** Sugar-free drinks (G2, Powerade Zero, Propel, Crystal Light or water) are also options.

A RESPONSIBLE DRIVER MUST REMAIN AT THE FACILITY THROUGHOUT YOUR APPOINTMENT. NO EXCEPTIONS.

Please read all instructions thoroughly.

If you are taking an injectable GLP medication for diabetes or weight loss you must not take the medication 7 days prior to your procedure.

** Please do not have nuts, seeds, or corn 3 days before your procedure**

1 Day BEFORE Your Exam:

Morning:

You will be on a clear liquid diet for 1 day before your procedure. This includes breakfast, lunch and dinner.
Clear liquids include water, sports drinks, black coffee, tea, any soda (light or dark), broth, gelatin dessert, ice pops and juice without pulp.
NOTHING RED, ORANGE OR PURPLE.

Noon:

- Take **Two (2) Dulcolax Tablets or Liquid Tablets** (bisacodyl laxative) by mouth.
- Mix one (1) bottle of MiraLAX with Gatorade in a separate container, and refrigerate it.
- If you can begin drinking the prep sooner, you may begin at 1PM – 2PM.

Evening:

At <u>4PM</u>, drink one (1) 8-ounce glass every 15 to 30 minutes until completed.



- You may slow down if you become nauseated.
- Continue drinking additional clear liquids throughout the evening.
- If you are not passing clear or yellowtinted water, mix the second MiraLAX bottle and Gatorade. Begin drinking the solution again until clear.
- If you are not passing clear or yellow-tinted water after drinking your prep, please call (318) 631-9121 for additional recommendations.

The Day of Your Exam:

- If you smoke, please do not smoke the day of your scheduled procedure.
- NOTHING BY MOUTH FOR 4 HOURS BEFORE YOU ARRIVE! (This includes water, gum and mints.)
 NOTHING by mouth; otherwise, your procedure may be canceled due to safety precautions.



Medications:

 Take your regularly scheduled blood pressure, heart and seizure medications with a sip of water at least two (2) hours before you arrive.



Diabetics:

- Test your blood sugar before coming in for your procedure, and bring the reading with you.
- If you take oral medication or insulin, **DO NOT** take it the morning of your procedure.
- Bring your insulin with you.
- Please contact the physician who manages your diabetic care if you have any questions regarding your oral medication or insulin dosages.



Blood Thinners:

- If you have not received instructions and are taking blood thinners (such as Coumadin, Warfarin or Plavix), call your prescribing physician, and ask if you may stop taking the medication 3 to 7 days prior to your procedure if you have not received instructions.
- You may continue taking aspirin, ibuprofen and other anti-inflammatory medications.



Bring with you:

- Completed paperwork forms that were sent to you
- Photo ID and Insurance Card
- List of all medications, vitamins and supplements you are taking (include dosage, and last date and time each medication was taken before your arrival)
- Wear a loose-fitting, short-sleeve shirt.

Is Your Bowel Clean?




- · Semi-solid stool
- Dark & murky





- Liquid stool
- Small amount of feces





- Liquid stool
- Particles present





- Liquid stool
- Particles present
- Almost ready





Clear liquid
READY

